

Jose Ubeda^{1,2,3,4}, Kenji Yoshikawa⁵, Walter Pari¹, David Palacios³, Pablo Masias¹, Fredy Apaza¹, Beto Ccallata¹, Rafael Miranda¹, Ronald Concha¹, Pool Vasquez¹ and Rolando Cruz²

¹Instituto Geológico Minero y Metalúrgico, Peru. ²Autoridad Nacional del Agua, Peru. ³Departamento de Análisis Geográfico Regional y Geografía Física. Universidad Complutense de Madrid, Spain. ⁴NGO Guías de Espeleología y Montaña, Spain. ⁵Water and Environmental Research Center. University of Alaska Fairbanks, USA. joseubeda@ucm.es

A network of air and ground temperature sensors installed 2004-2014 has enabled the discovery of permafrost on the Coropuna (6377 m) and Chachani (6057 m) volcanoes. However, on the Misti (5820 m) volcano there is no permafrost, which can be attributed to geothermal heat.

Misti and Chachani are very close to each other, near the city of Arequipa (S. Peru). Coropuna is 150 km to the west. Various volcanic eruptions have taken place on Misti and Coropuna in the last 10 ka (Úbeda et al, 2012).

The volcanic activity on the Chachani seems to be much older, although it has not been researched to date. Coropuna is covered by a glacial system of 40 km² (23-11-2013) and the moraines surrounding the volcanic complex indicate a surface of >500 km² >10 ka ago (Úbeda et al, 2011).

On Chachani the evidence also suggests a great extent in the past although in this case there are no glaciers conserved at the present day. On Misti there are currently no glaciers either, nor is there any evidence conserved of their earlier presence, and this has also been related to geothermal heat.

As well as other study areas, the CRYOPERU sensor network includes 4 stations in the sector Coropuna-NE; 3 stations in Coropuna-SE; 3 stations in Chachani-SE and 3 stations in Misti-NW.

The stations are at different altitudes, in an interval of 4300-6000 m. Each station has a thermometer to measure the air temperature (at a height of 0.50 m) and three thermometers to measure the ground temperature (at depths of 0.15, 0.30 and 1.00 m). The sensors are synchronized in GPS time and record the temperature every 30 minutes.

References

Úbeda, J. et al (2012). Glacial and volcanic evolution on Nevado Coropuna (Tropical Andes) based on cosmogenic ³⁶Cl surface exposure dating. EGU2012-3683-2.
Úbeda, J. (2011). El impacto del cambio climático en los glaciares del complejo volcánico Nevado Coropuna (Cordillera Occidental de los Andes Centrales). PhD Thesis. Universidad Complutense de Madrid. 594 pp. <http://eprints.ucm.es/12076/>

This research has been supported by the projects CRYOCRISIS CGL2012-35858 and www.cryoperu.pe.






Figure 1: General location on the Andean altiplano



Permafrost site Temperature stations

Figure 2: Temperature stations on Chachani



Permafrost site Temperature stations

Figure 3: Temperature stations on Coropuna




Figure 4: Fieldwork on Coropuna




Figure 5: Geophysical profiles in Coropuna



Figure 6: Permafrost from Coropuna




Figure 7: Geophysical profiles on Chachani




Figure 8: Soil temperatures on Chachani (monthly averages)



Figure 9: Permafrost from Chachani